

Director, Tamachi IC Clinic, Japan

2003 Graduated from Nihon University School of Medicine

Dr. Noguchi went to a drug rehabilitation facility in Australia during the medical internship. He learned the true cause of diseases and the importance of immunity and nutrition there. After returning to Japan, he has been working on comprehensive medical care including cancer care at private hospitals and clinics within the Tokyo metropolitan area. Meanwhile, he has developed various lecture activities and video lectures on the theme of preventive medicine and self-care, industrial physician activities at a special elderly nursing home and volunteer activities in the disaster area, etc.

The case of ovarian cancer in which the metastatic colon tumor resolved with combination use of BioBran (RBAC)-containing supplement and allogeneic leukocyte infusion therapy and avoided creation of an artificial anus

This case is a woman in her 40s. She had primary left ovarian cancer with the metastatic colon tumor. The doctor in charge told again and again her "You will have an artificial anus." But the metastatic colon tumor disappeared after combination use of BioBran (RBAC)-containing supplement and allogeneic leukocyte infusion therapy at Tamachi IC Clinic. And then she has succeeded in avoiding creation of the artificial anus.

The circumstances are as follows: When the examination laparotomy was performed under the doctor in charge, it was suggested that left ovarian cancer may have metastasized to the upper abdomen and rectum, and it was diagnosed as Stage III. The doctor suggested a policy to perform surgery including creation of an artificial anus after regression of the cancer by combination chemotherapy with Paclitaxel and Carboplatin. She felt anxiety about the treatment policy and the side effects of the chemotherapy, and she visited our clinic as a second opinion. On inquiry, it was discovered that she had constipation to the extent that she could not stop taking magnesium-containing preparations and Sennoside, decreased body temperature, and allergies to animals and house dust. On immunological examination, it was revealed that the activity of her natural killer cells remarkably decreased and the proportion of her lymphocytes decreased.

Therefore, for the purpose of supporting treatment to be carried out under the doctor in charge, our clinic recommended her to improve her lifestyle centered on improving eating habits, leukocyte infusion separated from allogeneic blood every about 2 weeks, and to take the supplement containing 0.2 g of BioBran (RBAC) per pack every day. But her constipation got worse after starting the chemotherapy. She also complained of abdominal pain, abdominal distension and anorexia. Thus, we recommended her to increase BioBran (RBAC)-containing supplements from 1 pack to 2 packs, 0.4 g of BioBran (RBAC) per day. As a result, after 2 weeks, her constipation was resolved and her abdominal distention got better. Her abdominal pain was also alleviated. Two weeks after that, her bowel movements were seen once a day without taking Sennoside. One month after that her appetite recovered. Four months after starting support of these, there was no malignant finding on the biopsy of the colonoscopy before the operation. Ultimately, the doctor in charge admitted the withdrawal of metastatic colon tumor, and then he operated her uterus and ovary only. That is to say, no artificial anus was constructed.